

Quantitative methylation detection in DNA samples

Field of the Invention

5

This invention relates to the analysis of nucleic acids, especially to the analysis of methylation patterns in genomic DNA by providing a means of detecting nucleotides, that are characteristic for methylated sites after bisulfite treatment of the genomic DNA. The method
10 utilises the incorporation of labels and the detection of fluorescence resonance energy transfer (FRET) of the amplified sample DNA.

15

Prior art

DNA Methylation

The levels of observation that have been studied in recent years in molecular biology have concentrated on
20 genes, the translation of those genes into RNA, and the transcription of the RNA into protein. There has been a more limited analysis of the regulatory mechanisms associated with gene control. Gene regulation, for example, at what stage of development of the individual a
25 gene is activated or inhibited, and the tissue specific nature of this regulation is less understood. However, it can be correlated with a high degree of probability to the extent and nature of methylation of the gene or genome. From this observation it is reasonable to infer
30 that pathogenic genetic disorders may be detected from irregular genetic methylation patterns.

The efforts of the Human Genome project are concentrated on the sequencing of the human genome. It is expected
35 that this will yield considerable therapeutic and diagnostic benefits for the treatment of disease.

However, these efforts have so far been unable to address a significant aspect of genetic disorders, the epigenetic factor. The epigenetic regulation of gene transcription has been shown to effect many disorders. One of the most significant epigenetic mechanisms so far identified has been the methylation of cytosine. The methylation of cytosine at the 5 position is the only known modification of genomic DNA. Although the exact mechanisms by which DNA methylation effects DNA transcription are unknown, the relationship between disease and methylation has been well documented. In particular methylation patterns of CpG islands within regulatory regions of genome appear to be highly tissue specific. Therefore, it follows that misregulation of genes may be predicted by comparing their methylation pattern with phenotypically 'normal' expression patterns. The following are cases of disease associated with modified methylation patterns.

- Head and neck cancer (Sanchez-Cespedes M et al. „Gene promoter hypermethylation in tumors and serum of head and neck cancer patients“ Cancer Res. 2000 Feb. 15;60 (4):892-5)
- Hodgkin's disease (Garcia JF et al „Loss of p16 protein expression associated wiht methylation of the p16INK4A gene is a frequent finding in Hodgkin's disease“ Lab invest 1999 Dec; 79 (12):1453-9)
- Gastric cancer (Yanagisawa Y et al. „ Methylation of the hMLH1 promoter in familial gastric cancer with microsatellite instability“ Int J Cancer 2000 Jan 1; 85 (1):50-3)
- Prader-Willi/Angelman's syndrome (Zeschnigh et al „Imprinted segments in the human genome: different DNA methylation patterns in the Prader Willi/Angelman syndrome region as determined by the genomic sequencing method“ Human Mol. Genetics (1997) (6) 3 pp 387-395)

- ICF syndrome (Tuck-Muller et al „CMDNA hypomethylation and unusual chromosome instability in cell lines from ICF syndrome patients" Cytogenet Cell Genet 2000; 89(1-2):121-8
- 5 - Dermatofibroma (Chen TC et al „Dermatofibroma is a clonal proliferative disease" J Cutan Pathol 2000 Jan;27 (1):36-9)
- Hypertension (Lee SD et al. „ Monoclonal endothelial cell proliferation is present in primary but not secondary pulmonary hypertension" J clin Invest 1998
- 10 Mar 1, 101 (5):927-34)
- Autism (Klauck SM et al. „Molecular genetic analysis of the FMR-1 gene in a large collection of autistic patients" Human Genet 1997 Aug; 100 (2) : 224-9)
- 15 - Fragile X Syndrome (Hornstra IK et al. „ High resolution methylation analysis of the FMR1 gene trinucleotide repeat region in fragile X syndrome" Hum Mol Genet 1993 Oct, 2(10):1659-65)
- Huntigton's disease (Ferluga J et al. „possible organ and age related epigenetic factors in Huntington's disease and colorectal carcinoma" Med hyptheses 1989
- 20 May;29(1);51-4

All of the above documents are hereby incorporated by

25 reference.

Bisulphite treatment

A relatively new and currently the most frequently used method for analyzing DNA for 5-methylcytosine is based

30 upon the specific reaction of bisulfite with cytosine which, upon subsequent alkaline hydrolysis, is converted to uracil which corresponds to thymidine in its base pairing behaviour. However, 5-methylcytosine remains unmodified under these conditions. Consequently, the

35 original DNA is converted in such a manner that methylcytosine, which originally could not be

distinguished from cytosine by its hybridisation behaviour, can now be detected as the only remaining cytosine using "normal" molecular biological techniques, for example, by amplification and hybridisation or sequencing. All of these techniques are based on base pairing which can now be fully exploited. In terms of sensitivity, the prior art is defined by a method which encloses the DNA to be analysed in an agarose matrix, thus preventing the diffusion and renaturation of the DNA (bisulfite only reacts with single-stranded DNA), and which replaces all precipitation and purification steps with fast dialysis (Olek A, Oswald J, Walter J. A modified and improved method for bisulphite based cytosine methylation analysis. Nucleic Acids Res. 1996 Dec 15;24(24):5064-6). Using this method, it is possible to analyse individual cells, which illustrates the potential of the method. However, currently only individual regions of a length of up to approximately 3000 base pairs are analysed, a global analysis of cells for thousands of possible methylation events is not possible. However, this method cannot reliably analyse very small fragments from small sample quantities either. These are lost through the matrix in spite of the diffusion protection.

An overview of the further known methods of detecting 5-methylcytosine may be gathered from the following review article: Rein, T., DePamphilis, M. L., Zorbas, H., Nucleic Acids Res. 1998, 26, 2255.

To date, barring few exceptions (e.g., Zeschnigk M, Lich C, Buiting K, Doerfler W, Horsthemke B. A single-tube PCR test for the diagnosis of Angelman and Prader-Willi syndrome based on allelic methylation differences at the SNRPN locus. Eur J Hum Genet. 1997 Mar-Apr;5(2):94-8) the bisulfite technique is only used in research. Always, however, short, specific fragments of a known gene are amplified subsequent to a bisulfite treatment and either

- completely sequenced (Olek A, Walter J. The pre-implantation ontogeny of the H19 methylation imprint. Nat Genet. 1997 Nov; 17(3):275-6) or individual cytosine positions are detected by a primer extension reaction
- 5 (Gonzalzo ML, Jones PA. Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). Nucleic Acids Res. 1997 Jun 15;25(12):2529-31, WO Patent 9500669) or by enzymatic digestion (Xiong Z, Laird PW. COBRA: a
- 10 sensitive and quantitative DNA methylation assay. Nucleic Acids Res. 1997 Jun 15;25(12):2532-4). In addition, detection by hybridisation has also been described (Olek et al., WO 99 28498).
- 15 Further publications dealing with the use of the bisulfite technique for methylation detection in individual genes are: Grigg G, Clark S. Sequencing 5-methylcytosine residues in genomic DNA. Bioessays. 1994 Jun;16(6):431-6, 431; Zeschnigk M, Schmitz B, Dittrich B,
- 20 Buiting K, Horsthemke B, Doerfler W. Imprinted segments in the human genome: different DNA methylation patterns in the Prader-Willi/Angelman syndrome region as determined by the genomic sequencing method. Hum Mol Genet. 1997 Mar;6(3):387-95; Feil R, Charlton J, Bird AP,
- 25 Walter J, Reik W. Methylation analysis on individual chromosomes: improved protocol for bisulphite genomic sequencing. Nucleic Acids Res. 1994 Feb 25;22(4):695-6; Martin V, Ribieras S, Song-Wang X, Rio MC, Dante R. Genomic sequencing indicates a correlation between DNA
- 30 hypomethylation in the 5' region of the pS2 gene and its expression in human breast cancer cell lines. Gene. 1995 May 19;157(1-2):261-4; WO 97 46705, WO 95 15373 and WO 45560.
- 35 Fluorescence Resonance Energy Transfer(FRET)

Fluorescence resonance energy transfer (FRET) is an interaction between two molecules wherein the excited state of one molecule (the donor) transfers energy to the other molecule (the acceptor). The donor molecule is a fluorophore while the acceptor molecule may or may not be. The energy transfer occurs without the emission of photons, and is based on dipole-dipole interactions between the two molecules. Molecules that are commonly used in FRET include fluorescein, N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX), 4-(4'-dimethylaminophenylazo) benzoic acid (DABCYL), and 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS). Basic conditions for FRET include the following:

- Close proximity between the donor and acceptor molecules (typically 10-100 Å).
- The emission spectrum of the donor molecule must overlap the absorption spectrum of the acceptor molecule. The transition dipole orientations of the donor and acceptor molecules must be approximately parallel.
- The extent of the energy transfer is dependent on the distance between the two molecules, and the overlap between the donor and acceptor spectra. It can be described by the following equation:

$$k_t(r) = t_{D-1} \cdot (R_0/r)^6$$

wherein r is the distance between the donor and the acceptor

t_D is the lifetime of the donor in the absence of energy transfer

R_0 is termed the Förster distance.

The efficiency of the energy transfer (for a single donor-acceptor pair) is given by:

$$E = R_0^6 / (R_0^6 + r^6)$$

Förster distances are typically in the range of 30-60 Å. Therefore FRET can be used as a highly sensitive method of measuring microscopic distances, this is particularly useful within the field of molecular biology where it has been utilised in a number of ways. It has been used in the study of protein structure, assembly, distribution, conformation and interactions, as well as the study of cell membranes and immunoassays. FRET has also been used in a number of ways in the analysis of nucleic acids. This includes the analysis of the structure and conformation of nucleic acids, hybridisation, PCR, sequencing and primer extension assays.

Another class of probe labels include fluorescence quenchers. The emission spectra of a quencher overlaps with a, fluorescent dye such that the fluorescence of the fluorescent dye is substantially diminished, or quenched, by the phenomena of fluorescence resonance energy transfer „FRET“ (Clegg (1992) Meth. Enzymol., 211:353-388). A fluorescent reporter dye and quencher joined in a configuration that permits energy transfer from the fluorophore to the quencher may result in a reduction of the fluorescence of the fluorescent dye. The reporter is a luminescent compound that can be excited either by chemical reaction, producing chemiluminescence, or by light adsorption, producing fluorescence. The quencher can interact with the reporter to alter its light emission, usually resulting in the decreased emission efficiency of the reporter. The efficiency of this quenching phenomenon is directly correlated with the distance between the reporter molecule and the quencher molecule (Yaron (1979) Analytical Biochemistry, 95:228-35).

Particular quenchers include but are not limited to rhodamine dyes such as tetramethyl-6-carboxyrhodamine

(TAMRA) or tetrapropano-6-carboxyrhodamine (ROX) (Bergot, U.S. Patent No. 5,366,860).

Enzymatic Amplification

5 PCR is a commonly used technique that has been described, for example in US patents numbers 4,683,195, 4,683,202, and 4,800,159. Briefly, it is the amplification of a nucleic acid sequence by repetitive cycles of annealing and extending primer to single stranded nucleic acids
10 followed by the denaturation of the resultant double stranded molecule. PCR, (and variations thereof) has a multitude of applications and is one of the key technologies involved in most forms of nucleic acid analysis and manipulation.

15 An important variation is the Multiplex-PCR, where more than 2 specific primer are used and a multitude of different specific amplifications are obtained in one reaction chamber.

20 There are several commonly used methods for the detection of PCR products, such as gel electrophoresis and the use of labelled primer oligonucleotides and nucleoside triphosphates. The use of fluorescent labelled
25 nucleotides and oligomers within PCR for nucleic acid analysis is also known.

PNA FRET probes

30 PNA can hybridise to its target complement in either a parallel or anti-parallel orientation. However, the anti-parallel duplex (where the carboxyl terminus of PNA is aligned with the 5' terminus of DNA, and the amino terminus is aligned with the 3'-terminus of DNA) is typically more stable (Egholm (1993) Nature, 365:566-68).
35 PNA probes are known to bind to target DNA sequences with high specificity and affinity (Coull, US Patent No.

6,110,676). The PNA FRET probe examples of the present invention, with reporter or quencher moieties, are designed such that the PNA anneals in the anti-parallel orientation with the target sequences.

5

PNA may be synthesized at any scale on automated synthesizers. The PNA FRET probes may be synthesized on many of the commonly used solid supports. After synthesis is complete, the PNA may be cleaved from the support,
10 purified, analysed and quantitated. Fluorescent-labeled PNA probes have demonstrated desirable properties in hybridization assays (Hyldig-Nielsen, U.S. Patent No. 5,985,563).

15 Genomic DNA for further amplification is obtained from DNA of cells, tissue or other test samples using standard methods. This standard methodology is found in references such as Fritsch and Maniatis eds., Molecular Cloning: A Laboratory Manual, 1989.

20

Real Time PCR

Real time PCR monitoring using fluorescence has been described in several manners. Firstly, the binding of double stranded DNA specific fluorescent dyes such as
25 ethidium bromide allows for the monitoring of the accumulation of PCR product by correlation with increased fluorescence. A second detection method, polymerase mediated exonuclease cleavage utilises the 5' exonuclease activity of polymerases such as Taq. An oligonucleotide
30 probe that is complementary to the PCR product, yet distinct from the PCR primer is labelled with a FRET pair such that the donor molecule is quenched by an acceptor molecule. During PCR amplification, the 5' exonuclease proceeds to digest the probe, separating the FRET pair
35 and leading to increased fluorescence. A variation on this technology uses a nucleic acid wherein the FRET pair

is internally quenched, for example, by having a hairpin conformation. Upon hybridisation to a sequence of interest, the FRET pair is separated and the donor molecule emits fluorescence. This technology can be used, for example for the analysis of SNPs.

An alternative technology is based on the use of two species of hybridisation probes, each labelled with a member of a FRET pair. Upon hybridisation of both probes to the target sequence in adequate proximity, a fluorescent signal is emitted. Again, this technology may be used for the detection of SNPs.

A major advantage of the use of such FRET based PCR technologies is that the reaction may be monitored in a closed tube reaction, suitable for use in high and medium throughput and reducing the probability of contamination.

Description of the Invention

According to the present invention there is provided a method for the cytosine methylation detection in a DNA sample, comprising the following steps:

- a) a genomic DNA sample is treated in a manner capable of distinguishing methylated from unmethylated cytosine bases;
- b) the pre-treated DNA is amplified using at least one oligonucleotide primer, a polymerase and a set of nucleotides of which at least one is marked with a first type of label;
- c) a sequence-specific oligonucleotide or oligomer probe is hybridized to the amplification product and a FRET occurs if the oligonucleotide or oligomer probe, marked with a second type of label, binds in close proximity to one of the labeled nucleotides that was incorporated into the amplification product;

d) the level of methylation of the sample is determined by the level of interaction between said first and second type of label.

5 According to the invention it is preferred that the first type of label is a donor fluorophore and the second type of label is an acceptor fluorophore and that the extent of fluorescence resonance energy transfer (FRET) is measured. It is further preferred that the first type of
10 label is an acceptor fluorophore and the second type of label is a donor fluorophore and that the extent of fluorescence resonance energy transfer (FRET) is measured.

15 A further preferred embodiment of the present invention is characterised in that the nucleotides of step b) contain a fluorescent moiety and the probe in step c) a quencher moiety. It is also preferred according to the invention that the nucleotides of step b) contain a
20 quencher moiety and the probe in step c) a fluorescent moiety.

According to the invention it is also preferred that the polymerase has no 5' to 3' exonuclease activity in order
25 to prevent degradation of the probe.

It is further preferred according to the invention that a change in fluorescence intensity is monitored in real-time during the amplification reaction.

30 It is also especially preferred according to the present invention that a change in fluorescence intensity is monitored at end-point of target amplification.

According to another preferred embodiment of the present invention the amplification reaction is achieved with the polymerase chain reaction (PCR).

5 According to the invention it is preferred that the probe contains only one CpG or that the probe contains several CpGs. Especially in this case it is further preferred that each probe for each CpG has a fluorescent label.

10 In a further preferred embodiment of the present invention the probe can be end labeled or internally labeled.

15 It is also preferred according to the invention that the methylation information is determined by the change in fluorescence intensity during subsequent rounds or cycles of PCR.

20 It is also preferred that the sample DNA is only amplified by chosen PCR primers if a certain methylation state is present at a specific site in the sample DNA.

25 According to the present invention a method is preferred wherein the sample DNA is only amplified if a certain methylation state was present at a specific site in the sample DNA, the sequence context of which is essentially complementary to one or more oligonucleotides or PNA oligomers which are additionally used in the PCR reaction.

30 It is also preferred that the amplification from the 3'-end of the probe is blocked by phosphorylation.

35 According to the invention it is also preferred that a melting curve is generated at the end of the PCR to gather additional data.

It is especially preferred within the scope of the present invention that the fluorescent moiety is a fluorescein dye, a rhodamine dye, or a cyanine dye. Especially preferred is also that the quencher moiety is a rhodamine dye.

It is an especially preferred feature of the present invention that the deamination treatment of the DNA is performed with a bisulfite reagent.

It is also preferred according to the invention that the DNA sample is cleaved prior to deamination treatment with restriction endonucleases.

In a preferred embodiment of the method of the invention the DNA sample is isolated from mammalian sources e.g. cell lines, blood, sputum, faeces, urine, cerebrospinal fluid, tissue embedded in paraffin, for example, ocular tissue, intestine, kidney, brain, heart, prostate, lung, chest or liver, histological slides and all possible combinations.

It is another preferred embodiment of the present invention to use of a pre-treated genomic DNA within the method according to the present invention for the determination of the methylation status of a corresponding genomic DNA.

Another object of the present invention is to provide a diagnostic kit for the detection of the methylation of cytosine bases in genomic DNA samples, comprising reagents for the selective deamination of cytosine bases in genomic DNA, one or more primers and labeled nucleotides for the amplification step, a detectable

probe and optionally protocols or instructions for one of the methods according to one of the preceding claims.

5 The invention describes a method to determine the presence of specific CpG dinucleotides in a fragment of DNA using fluorescence resonance energy transfer (FRET). This can be used to obtain information about sequence properties of a sample DNA fragment. For example, a point mutation could be identified in a fragment if a
10 nucleotide is present in its sequence as a result of this mutation which is not present in the wild type.

15 The method is preferably used to measure cytosine methylation. As mentioned above, bisulphite leads to the selective deamination of cytosine, leaving 5-methylcytosine essentially unchanged. Methylation of cytosine occurs almost exclusively in the sequence context 5'-CG-3'. Therefore, after bisulphite treatment, certain dinucleotides containing C do not occur anymore
20 in one strand, but they may still occur in the complementary strand formed in the amplification of bisulphite treated DNA, for example using the polymerase chain reaction (PCR).

25 This invention provides a method for visualizing the methylation status of a CpG at defined positions in a very sensitive way with very low background signal.

30 The method briefly comprises the following steps of

- a) treating a genomic DNA sample in a manner capable of distinguishing methylated from unmethylated cytosine bases;
- 35 b) amplifying the pre-treated DNA using at least one oligonucleotide primer, a polymerase and a set of

nucleotides of which at least one is marked with a first type of label;

5 c) hybridising a sequence-specific oligonucleotide or oligomer probe to the amplification product, a FRET occurs if the oligonucleotide or oligomer probe, marked with a second type of label, binds in close proximity to one of the labeled nucleotides that was incorporated into the amplification product;

10

d) determining the level of methylation of the sample by measuring by the level of interaction between said first and second type of label.

15

In a preferred embodiment of the invention, the first type of label is a donor fluorophore and the second type of label is an acceptor fluorophore and that the extent of fluorescence resonance energy transfer (FRET) is measured.

20

In a further preferred embodiment of the invention, said first type of label is an acceptor fluorophore and the second type of label is a donor fluorophore and that the extent of fluorescence resonance energy transfer (FRET) is measured.

25

It is preferred that two or more CpGs are separately interrogated in an amplification reaction.

30

Preferably, separate probes for each CpG are used, each with its own fluorescent label.

35

The instant invention also enables a multiplex PCR to rapidly determine optimal assay parameters and a fast, cost-effective, and accurate system for the quantitative analysis of target analytes. A multiplexed assay can for

example be designed in a standard 96 well microtiter plate format at room temperature using conventional robotic systems for sample delivery and preparation.

- 5 Preferably, the oligonucleotide or oligomer probes used comprise one or more nucleotide analogs selected from a nucleobase analog, a 2'-deoxyribose analog, an internucleotide analog, PNA or LNA.
- 10 Although the detection of C or G nucleotides (in the complementary strand) after bisulphite treatment in the context CpG is the preferred application, also any other nucleotide or polymorphism can in principle be detected. The preferred application is to determine the methylation
- 15 status of certain CpG positions by determining the level of interaction between an unconverted C in the bisulphite treated DNA and a labeled oligonucleotide hybridised thereto. Accordingly, Guanin in the complementary strand (after PCR) can be used for the same purpose.
- 20 Also, converted positions after bisulphite treatment can be identified by detecting thymine (or adenine in the complementary strand) at selected positions using this technology. However, design of probes becomes more
- 25 difficult as it is not possible to distinguish between T that was in the original genomic sequence and T positions that were created through bisulphite conversion, indicating lack of methylation at the respective cytosines.
- 30 In more detail, this method for the detection of specific nucleotides in a DNA sample is characterized in that an isolated genomic DNA sample is treated in a manner capable of distinguishing methylated from unmethylated
- 35 cytosine bases and the pre-treated DNA is amplified using at least one oligonucleotide primer, a polymerase and a

set of nucleotides at least one of which is marked with a first type of label, in a first embodiment a donor fluorophore and in a second embodiment an acceptor fluorophore.

5

Preferably, the acceptor and donor dyes (fluorophores) are chosen in a way that the emission wavelength of the donor dye overlaps with the excitation wavelength of the acceptor dye. It is preferred that the emission and excitation spectra are sharp peaks and that the emission spectra of the dyes are unlikely to overlap.

10

It is preferred that the polymerase has no 5' to 3' exonuclease activity to prevent degradation of the probe.

15

For example, dGTP is labeled with a fluorescent dye. The labeled dGTP is incorporated only where there was a methylated cytosine in the original DNA sample. Alternatively, the dCTP can be labeled with a fluorescent dye.

20

After the extension phase, the DNA is denaturated and then allowed to reanneal.

25

A sequence-specific oligonucleotide or oligomer probe (referred to as oligomer probes if DNA analogs like PNA are used), marked with a second type of label that is in a first embodiment an acceptor fluorophore and in a second embodiment a donor fluorophore hybridizes to the amplification product. Preferably, amplification from the 3' end of the probe is blocked by phosphorylation (with dideoxynucleotides).

30

The labels are preferably introduced into the oligonucleotide probes by standard enzymatic methods, such as the use of 5' labeled amplification primers for

35

5' labeling or fluorescent-labeled base analogs for internal labeling.

A FRET reaction occurs if the fluorescently labeled oligonucleotide, preferably in one embodiment a fluorescent moiety binds in close proximity to a nucleotide labeled with a quencher moiety, that was incorporated into the amplification product or in another embodiment a quencher moiety that binds in close proximity to a nucleotide labeled with a fluorescent moiety that was incorporated into the amplification product (Fig.1).

It is preferred that the probe is positioned in several places relative to the CpG. The conformation of the DNA controls the spacing of the two fluorescent dyes. Preferently, the positioning is optimized for each CpG.

In a preferred embodiment, the probe is separated from the amplification primers or alternatively it is attached to the 5' end of one of the primers.

This way, the level of methylation can be determined, identifying the CG dinucleotides. If a TG dinucleotide is present instead, no FRET will be observed. Therefore, this method can be directly used to monitor DNA methylation (Fig. 2).

In a preferred embodiment of the invention a real time monitoring of the FRET signal is performed during the amplification reaction. This way, the progress of the amplification can be examined (Fig.3). Very preferably the amplification reaction is a polymerase chain reaction (PCR), even though other amplification procedures for example cloning or SDA (Strand Displacement Amplification) are also preferred.

In another preferred embodiment of the invention the change in fluorescence intensity is monitored at end-point of target amplification. End-point analysis of the PCR entails fluorescent dye signal measurement when thermal cycling and amplification is complete. Results are reported in terms of the change in fluorescence, i.e. fluorescence intensity units, of the fluorescent dye signal from start to finish of the PCR thermal cycling, preferably minus any internal control signals.

It is also preferred that a melting curve is generated at the end of the PCR to gather additional data.

In another preferred embodiment of the invention the CpG dinucleotide occurs only once in the amplification product. As outlined above, this is very helpful if the presence of the FRET signal is directly used to draw conclusions about the sequence characteristics of the sample DNA.

For example, if only one labeled CG is present in an amplification product of a bisulphite treated sample and a fluorescently labeled probe binds in close proximity to it, a FRET occurs and direct conclusions can be drawn that a methylated cytosine was present in a certain position in the genomic DNA sample.

If several labeled CGs are present for example in an amplification product of a bisulphite treated sample and probes bind in close proximity to them, each with its own fluorescent label, several FRET reactions occur and conclusions can be drawn about the methylated cytosines from all sites involved.

In a further preferred embodiment of the invention the methylation information is determined by the change in fluorescence intensity during subsequent rounds of PCR.

- 5 Preferably, the sample is illuminated during the amplification reaction with light of appropriate wavelength.

10 In a preferred embodiment of the invention, prior to the PCR either essentially all cytosines in the DNA sample are selectively deaminated, but 5-methylcytosines remain essentially unchanged or essentially all 5-methylcytosines in the DNA sample are selectively deaminated, but cytosines remain essentially unchanged.

15 Cytosine-guanine (CpG) dinucleotides are detected, allowing conclusions about the methylation state of cytosines in said CpG dinucleotides in said DNA sample. This deamination is preferably performed using a bisulphite reagent.

20 Preferably, the sample DNA is only amplified by chosen PCR primers if a certain methylation state is present at a specific site in the sample DNA the sequence context of which is essentially complementary to one or more of said

25 chosen PCR primers. This can be done using primers annealing selectively to bisulphite treated DNA which contains in a certain position either a TG or a CG, depending on the methylation status in the genomic DNA. Primers can be designed for both cases. A primer could

30 contain a G at its 3'-end, therefore it would only bind to a DNA containing a C at the respective position and therefore this primer will only or preferentially amplify methylated DNA because the C is indicative of a methylation in this position after bisulphite treatment.

35 This method is known as MSP, methylation sensitive PCR.

In another preferred embodiment of the invention, the sample DNA is only amplified if a certain methylation state is present at a specific site in the sample DNA the sequence context of which is essentially complementary to one or more oligonucleotides or PNA oligomers which are additionally used in the PCR reaction. These oligonucleotides or PNA oligomers bind selectively to the template DNA and prevent its amplification depending on the methylation state of the DNA prior to bisulphite conversion.

Preferably, the fluorescent moiety is a fluorescein dye, a rhodamine dye, or a cyanine dye and the quencher moiety a rhodamine dye.

In another preferred variant of the invention the DNA sample is cleaved prior to deamination (for example bisulphite) treatment with restriction endonucleases.

Preferred is also a method whereby the enzymatic amplification of the treated DNA is such that only one strand of the DNA sample is amplified.

Preferably, the DNA sample is isolated from mammalian sources e.g. cell lines, blood, sputum, faeces, urine, cerebrospinal fluid, tissue embedded in paraffin, for example, ocular tissue, intestine, kidney, brain, heart, prostate, lung, chest or liver, histological slides and all possible combinations.

Another embodiment of the present invention is a diagnostic kit for the detection of the methylation of cytosine bases in genomic DNA samples, comprising reagents for the selective deamination of cytosine bases in genomic DNA, one or more primers and fluorescently labeled nucleotides for the amplification step and

optionally protocols or instructions for one of the methods according to one of the preceding claims.

5 This kit can also comprise several additional items for example detectable probes.

10 The components of said kit, as an example, could comprise receptacles for the following in sufficient quantities to carry out the method:

- 1) Reagents for the bisulfite conversion of sample DNA.
- 2) Reagents for the amplification of the converted sample and incorporation of fluorophore labelled nucleotides including:
 - 15 a) nucleic acid primer and
 - b) appropriate mix of nonlabeled and fluorophore labeled nucleotides and
 - c) DNA polymerase able to incorporate the fluorophore labelled nucleotides
- 20 3) Instructions for use

25 The term 'instructions for use' should cover a tangible expression describing the reagent concentrations for the assay method, parameters such as the relative amounts of reagents to be combined, maintenance times for reagents/sample mixtures, temperature, buffer conditions and the like.

30 In the following, steps of preferred embodiments of the invention are described in more detail.

DNA Isolation

35 The genomic DNA sample must be isolated from tissue or cellular sources. For mammals, more preferably humans, the DNA sample may be taken from any tissue suspected of expressing the target site within the genome and also

from, such as cell lines, blood, sputum, faeces, urine, cerebrospinal fluid, tissue embedded in paraffin; for example, tissue of intestine, kidney, brain, heart, prostate, lung, chest or liver, histological slides, but not limited to those. Extraction may be by means that are standard to one skilled in the art, these include the use of detergent lysates, sonification and vortexing with glass beads. Once the nucleic acids have been extracted the genomic double stranded DNA is used for analysis.

10

Bisulfite Treatment

The sample DNA is then treated chemically in order to convert the unmethylated cytosine bases into uracil. The chemical modification may be by means of, for example, (but not limited to) a bisulfite solution. Said chemical conversion may take place in any format standard in the the art. This includes but is not limited to modification within agarose gel or in denaturing solvents.

Wherein the chemical modification takes the form of a bisulfite treatment of the DNA the following steps may be followed.

The double stranded DNA must be denatured. This may take the form of a heat denaturation carried out at variable temperatures. For high molecular weight DNA, the denaturation temperature is generally greater than 90 °C. However, the analysis may be upon smaller fragments which do not require such high temperatures. In addition as the reaction proceeds and the cytosine residues are converted to uracil the complementarity between the strands decreases. Therefore, a cyclic reaction protocol may consist of variable denaturation temperatures.

The bisulfite conversion then consists of two important steps, the sulfonation of the cytosine and the subsequent

deamination. The equilibra of the reaction are on the correct side at two different temperatures for each stage of the reaction. Taking into account the kinetics of the reactions it is preferable that the reaction takes place under cyclic conditions, with changing temperatures. The temperatures and length at which each stage is carried out may be varied according to the specific requirement of the situation. However, a preferred variant of the method comprises a change of temperature from 4 C (10 minutes) to 50 C (20 minutes). This form of bisulfite treatment is state of the art with reference to WO 99/28498.

Said chemical conversion may take place in any format standard in the art. This includes but is not limited to modification within agarose gel, in denaturing solvents or within capillaries.

Bisulfite conversion within agarose gel is state of the art and has been described by Olek et al, Nucl. Acids. Res. 1996, 24, 5064-5066. The DNA fragment is embedded in agarose gel and the conversion of cytosine to uracil takes place with hydrogensulfite and a radical scavenger. The DNA may then be amplified without need for further purification steps.

In a further preferred embodiment the DNA conversion may take place without an agarose matrix. The DNA may incubated at increased temperatures with hydrogensulfite and a radical scavenger. Said reaction takes place within an organic denaturing solvent. Examples of denaturing solvents include, but are not limited to, Polyethylene glycol dialkyl polyethylenglycol-dialkylether, dioxane and substituted derivatives, urea or derivatives, acetonitrile, primary alcohols, secondary alcohols, tertiary alcohols, DMSO or THF.

In a further embodiment, prior to chemical treatment the DNA sample is transferred into a heatable capillary that is permeable to small molecules. The reaction steps of the chemical modification may then be carried out in the capillary tubes by means of the addition and removal of reagents through connected capillaries.

Subsequent to the chemical treatment the two strands of the DNA may no longer be complementary.

Amplification and Incorporation of labeled Nucleotides
Fractions of the so treated genomic DNA are then enzymatically amplified using oligonucleotide primers. The length and design of said primers may be specific to the area of the genome to be analysed. As such a wide range of primers are suitable for use in this technique. Such primer design is within the state of the art.

An appropriate fraction of the nucleotides presented in the amplification reaction, for example the G nucleotides, are labeled with either a first or second type of label, the first type being a fluorophore and the second type a quencher or vice versa. Acceptable fluorophores for labeling the nucleotides are well known to those skilled in the art and include, but are not limited to, fluorescein, rhodamine, cyanine, phycoerythrin, Cy 5, Cy 5.5, Cy 7, LC Red 640 or LC Red 705 whereas the acceptable quenchers are rhodamine dyes. Attaching those dyes to the nucleotides lies within the state of the art.

In a preferred embodiment of this invention the sample is illuminated during the amplification reaction with light of appropriate wavelength.

The skill of the invention lies in the interpretation of a FRET signal during states, where a sequence-specific oligonucleotide probe is hybridized to an amplification product and a FRET occurs if the fluorescently labeled oligonucleotide binds in close proximity to one of the labeled nucleotides that were incorporated into the amplification product in order to gain knowledge of the methylation state of the sample.

Advanced Data Processing

It is anticipated that the method will be used for the high throughput analysis of genomic DNA samples. Therefore, the invention also involves analysis of data using a computing device. In a preferred embodiment said device may comprise one or more databases. In a further preferred embodiment said device may comprise one or more learning algorithms.

Description of the Drawings

Figure 1

Legend:

- 1: Bisulfite Treatment
- 2: Incorporation of labeled nucleotide
- 3: Detection of target by Fret
- 4: Real-Time PCR

DNA extracted from a tissue is treated with sodium bisulfite (A). Into the bisulfite treated DNA (single strand (B) dGTPs are incorporated (C+D) that are labeled with a fluorescent dye during a real-time PCR. A FRET occurs if the fluorescently labeled oligonucleotide binds in close proximity to one of the labeled nucleotides that was incorporated into the amplification product (E).

Figure 2

Legend:

1. Bisulfite treatment
- 5 2. PCR with labeled dGTP
3. Extension of primer
4. Denaturation
5. Annealing and fluorescence monitoring
6. More Cycles

10

15

20

DNA of interest is chemically treated, the only cytosines remaining in the sequence are those that were methylated in the original sample. PCR primers designed to target one of the DNA strands anneal to the template and extend it by incorporating labeled nucleotides (dGTP). The labeled dGTP is incorporated only where there was a methylated cytosine in the original DNA sample. After the extension phase, the DNA is denaturated, allowed to reanneal and the fluorescence monitored. With each round of PCR, more targets complementary to the probe accumulate. The amount of fluorescence emitted from the probe is measured.

Figure 3

25

Legend:

1. PCR with unlabeled dCTP, dATP, dTTP and labeled dGTP
2. Hybridization of fluorescent-labeled gene specific oligo-nucleotide
- 30 3. FRET, real-time fluorescence detection

35

A PCR is performed with primers that target one of the DNA strands. The first primer anneals to the template and extends it by incorporating the appropriate nucleotides. One of the nucleotides, in this case dGTP, is labeled with a fluorescent dye. A sequence-specific

oligonucleotide probe hybridizes to the site of interest.
If the labeled guanine is present, a FRET reaction
occurs. The energy emitted from the guanine is
transferred to the label on the probe. The energy emitted
5 from the probe is detected by real-time fluorescence
detection.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221